

using more recent data is unlikely to change that conclusion (see Appendix of Jerrett et al. 2009). Finally, national risk estimates are more applicable globally than city-specific estimates because they include larger and more diverse populations.

However, because the evidence for chronic ozone mortality is more limited than the large body of evidence demonstrating mortality associations with short-term ozone exposure, we present here estimates of the global burden of ozone on mortality using RR estimates from Bell et al. (2004), a large multicity study of short-term ozone mortality. We estimated mortalities daily using the difference between preindustrial and present-day 8-hr maximum ozone, and sum mortalities over the 1-year simulation. We used the reported relationship for cardiopulmonary mortality and daily average ozone [0.64% (95% posterior interval, 0.31–0.98%) for a 10-ppb increase], and corrected to 8-hr ozone using the reported ratio between daily 8-hr and 24-hr average ozone associations with nonaccidental mortality.

Using these methods, we estimated 362,000 (95% confidence interval, 173,000–551,000) annual global premature cardiopulmonary deaths attributable to ozone, approximately 50% of the 700,000 premature deaths we calculated in our original study (Anenberg et al. 2010). Since estimated deaths due to PM_{2.5} (3.7 million) are an order of magnitude larger, using a short-term rather than long-term RR estimate for ozone has only a minor effect on the overall global burden of disease due to outdoor air pollution. As RRs for chronic ozone mortality are not as strongly supported as those for PM_{2.5}, we expect that estimates of mortality burden will improve as research on chronic ozone exposure and mortality continues globally.

S.C.A. and J.J.W. received unrestricted funding from the Merck Foundation to support a related project, and the Merck Foundation was unaware of this research. D.Q.T. is an employee of Science and Technology Corporation, working at NOAA facilities. L.W.H. declares he has no competing financial interests.

Susan C. Anenberg
J. Jason West

The University of North Carolina
at Chapel Hill
Chapel Hill, North Carolina
E-mail: jasonwest@unc.edu

Larry W. Horowitz
NOAA Geophysical Fluid
Dynamics Laboratory
Princeton, New Jersey

Daniel Q. Tong
Science and Technology Corporation at
NOAA Air Resources Laboratory
Silver Spring, Maryland

REFERENCES

- Anenberg SC, Horowitz LW, Tong DQ, West JJ. 2010. An estimate of the global burden of anthropogenic ozone and fine particulate matter on premature human mortality using atmospheric modeling. *Environ Health Perspect* 118:1189–1195.
- Bell ML, McDermott A, Zeger SL, Samet JM, Dominici F. 2004. Ozone and short-term mortality in 95 US urban communities, 1987–2000. *JAMA* 292:2372–2378.
- Jerrett M, Burnett RT, Pope CA III, Ito K, Thurston G, Krewski D, et al. 2009. Long-term ozone exposure and mortality. *N Engl J Med* 360:1085–1095.
- Krewski D, Jerrett M, Burnett RT, Ma R, Hughes E, Shi Y, et al. 2009. Extended Follow-up and Spatial Analysis of the American Cancer Society Study Linking Particulate Air Pollution and Mortality. Research Report 140. Boston:Health Effects Institute.
- National Research Council, Committee on Estimating Mortality Risk Reduction Benefits from Decreasing Tropospheric Ozone Exposure. 2008. Estimating Mortality Risk Reduction and Economic Benefits from Controlling Ozone Air Pollution. Washington, DC:National Academies Press. Available: http://www.nap.edu/catalog.php?record_id=12198#toc [accessed 11 March 2011].

Influence of Selenium and Mercury on Age-Related Cataracts in the Brazilian Amazon

doi:10.1289/ehp.1003242

In their article, Lemire et al. (2010) provided important data on the frequency of age-related cataracts among adults in the Brazilian Amazon and found a correlation between age-related cataracts and whole-blood total mercury (Hg) concentrations and selenium (Se) levels in plasma and whole blood. However, in the “Discussion” of their paper, they stated that they “observed no adverse effects although Se concentrations were very high, reaching 1,500 µg/L for [blood]-Se and 913 µg/L for [plasma]-Se” (Lemire et al. 2010). However, they did not mention the potential and substantial adverse health effects associated with a high body Se burden. There are potentially adverse consequences to Se body burden, such as hair loss (alopecia), tooth decay, nail changes, peripheral paresthesias, weakness, skin lesions, and diabetes (Hira et al. 2004; Nuttall 2006; Shearer 1975; Stranges et al. 2010; Sutter et al. 2008; Yang et al. 1983). It would be valuable to know what Se-related adverse effects were observed by Lemire et al. (2010). Most studies, taken together, suggest a possible attenuation of Hg toxicity, probably as insoluble form of Hg selenide (Clarkson 2002; International Programme on Chemical Safety 1990).

Selenium may be able to delay the onset of toxic effects in animal models exposed to methylmercury in the diet (Clarkson 2002; Ganther et al. 1972). However, the Hg–Se interaction may not have an equivalent effect in some animals. There is evidence that coadministration of methylmercury and Se may lead to an important synergistic effect (Heinz and Hoffman 1998). In studies of persons who have been co-exposed to Hg

and Se, toxic effects of Se should be taken into account to discover the potential synergistic effect between Se and Hg.

The authors declare they have no competing financial interests.

Claudio Minoia
Anna Ronchi

Laboratory of Environmental and
Toxicology Testing
“S. Maugeri” IRCCS
Pavia, Italy

Paolo D. Pigatto

Department of Technology for Health
Dermatological Clinic
IRCCS Galeazzi Hospital
University of Milan
Milan, Italy

Gianpaolo Guzzi

Italian Association for Metals and
Biocompatibility Research–AIRMED
Milan, Italy
E-mail: gianpaolo_guzzi@fastwebnet.it

REFERENCES

- Clarkson TW. 2002. The three modern faces of mercury. *Environ Health Perspect* 110(suppl 1):11–23.
- Ganther HE, Goudie C, Sunde ML, Kopecky MJ, Wagner P. 1972. Selenium: relation to decreased toxicity of methylmercury added to diets containing tuna. *Science* 175:1122–1124.
- Heinz GH, Hoffman DJ. 1998. Methylmercury chloride and selenomethionine interactions on health and reproduction in mallards. *Environ Toxicol Chem* 17:139–145.
- Hira CK, Partal K, Dhillon KS. 2004. Dietary selenium intake by men and women in high and low selenium areas of Punjab. *Public Health Nutr* 7:39–43.
- International Programme on Chemical Safety. 1990. Methylmercury. *Environmental Health Criteria* 101. Geneva:International Programme on Chemical Safety, World Health Organization. Available: <http://www.inchem.org/documents/ehc/ehc/ehc101.htm> [accessed 2 March 2011].
- Lemire M, Fillion M, Frenette B, Mayer A, Philibert A, Passos CJ, et al. 2010. Selenium and mercury in the Brazilian Amazon: opposing influences on age-related cataracts. *Environ Health Perspect* 118:1584–1589.
- Nuttall KL. 2006. Evaluating selenium poisoning. *Ann Clin Lab Sci* 36:409–420.
- Shearer TR. 1975. Developmental and postdevelopmental uptake of dietary organic and inorganic selenium into the molar teeth of rats. *J Nutr* 105:338–347.
- Stranges S, Laclaustra M, Ji C, Cappuccino FP, Navas-Acien A, Ordoñez JM, et al. 2010. Higher selenium status is associated with adverse blood lipid profile in British adults. *J Nutr* 140:81–87.
- Sutter ME, Thomas JD, Brown J, Morgan B. 2008. Selenium toxicity: a case of selenosis caused by a nutritional supplement. *Ann Intern Med* 148:970–971.
- Yang GQ, Wang SZ, Zhou RH, Sun SZ. 1983. Endemic selenium intoxication of humans in China. *Am J Clin Nutr* 37:872–881.

Influence of Selenium: Lemire et al. Respond

doi:10.1289/ehp.1003242R

We agree with the comments made by Minoia et al. in their letter. In this study population, we did evaluate the sentinel signs and symptoms of selenosis (Lemire M, Philibert A, Fillion M, Passos CJS, Guimarães JRD,

Barbosa F Jr, Mergler D, unpublished data), and we observed no association between any of the biomarkers of selenium (Se) status and signs and symptoms of selenosis (hair loss, broken nail walls, nail sloughing, skin lesions, garlic breath, gastrointestinal disorders, and motor and sensory deficits), despite high Se body burdens. Other results in this study population show a positive association between Se status and motor performance (Lemire M, Fillion M, Frenette B, Passos CJS, Guimarães JRD, Barbosa F Jr, Mergler D, unpublished data).

Guidelines on Se toxicity (International Programme on Chemical Safety 1986; U.S. Environmental Protection Agency 2002) are based mostly on reports of chronic selenosis in a Chinese population with excessive Se exposure, resulting from high Se in crops fertilized with coal ash highly rich in Se (Yang et al. 1983), and from combustion of Se-rich coal for domestic use (Liu et al. 2007). Drinking water likewise contained unusually high concentrations of inorganic Se (Yang et al. 1983). Thus, several factors, such as exposure to toxic vapors from coal combustion and/or inorganic Se from drinking water, may have contributed to toxic Se effects observed in China, primarily attributed to high organic Se in local crops. Recent studies reporting selenosis most often refer to excessive Se intake from nutritional supplements or inorganic Se through drinking water (Sutter et al. 2008; Vinceti et al. 2010).

Our study (Lemire et al. 2010) was conducted in 2006, before the first publication on an association between Se supplementation and the incidence of self-reported type 2 diabetes (Stranges et al. 2007). In the Amazonian population in our study, only 5 of the 448 participants reported diabetes. Further studies should address the possible associations between high Se status and hypertension and hypercholesterolemia (reviewed by Stranges et al. 2010).

Other evidence comes from Inuit, whose traditional diet of marine mammals is exceptionally rich in Se. The prevalence of diabetes in Nunavik, Québec, Canada, is low (3.5%). Ferland et al. (2009) observed no association between Se status and diabetes or plasma fasting glucose and insulin levels. Inverse association between blood Se and systolic blood pressure has also been reported (Valera et al. 2009). Hansen et al. (2004) pointed out that there are no recorded signs of selenosis in Greenland populations and that their high Se intake may be tolerated at higher levels.

In Northern and Amazonian populations exposed to mercury (Hg), high dietary Se may offset Hg-mediated oxidative stress and/or be required to maintain optimal selenoenzymes. There may be consequently less

“excess” Se and little or no Se toxicity (Khan and Wang 2009).

Whether dietary Se is less toxic in Hg-exposed populations remains unanswered. However, even if there is increasing evidence that Se can offset some toxic effects of Hg, it may be inefficient against all Hg-mediated effects. Preventive actions should continue to focus on reducing Hg exposure rather than increasing Se status.

The authors declare they have no competing financial interests.

Mélanie Lemire

Axe santé des populations
et environnementale

Centre de recherche du Centre hospitalier
universitaire de Québec
Québec, Québec, Canada

E-mail: Melanie.Lemire@crchuq.ulaval.ca

Donna Mergler

Centre de recherche interdisciplinaire
sur la biologie, la santé, la société
et l'environnement

Université du Québec à Montréal
Montréal, Québec, Canada

REFERENCES

- Ferland A, Chateau-Degat ML, Lamarche B, Counil E, Dewailly É. 2009. Serum and toenail selenium and type 2 diabetes: learnings from a population with high dietary selenium intakes [Abstract]. *Diabetes* 58(suppl 1):A587. Available: http://professional.diabetes.org/Abstracts_Display.aspx?TYP=1&CID=72799 [accessed 3 March 2011].
- Hansen JC, Deutch B, Pedersen HS. 2004. Selenium status in Greenland Inuit. *Sci Total Environ* 331(1–3):207–214.
- International Programme on Chemical Safety. 1986. Selenium. *Environmental Health Criteria* 58. Geneva:International Programme on Chemical Safety, World Health Organization. Available: <http://www.inchem.org/documents/ehc/ehc/ehc58.htm> [accessed 3 March 2011].
- Khan MAK, Wang F. 2009. Mercury-selenium compounds and their toxicological significance: toward a molecular understanding of the mercury-selenium antagonism. *Environ Toxicol Chem* 28(8):1567–1577.

- Lemire M, Fillion M, Frenette B, Mayer A, Philibert A, Passos CJ, et al. 2010. Selenium and mercury in the Brazilian Amazon: opposing influences on age-related cataracts. *Environ Health Perspect* 118:1584–1589.
- Liu GJ, Zheng LG, Duzgoren-Aydin NS, Gao LF, Liu JH, Peng ZC. 2007. Health effects of arsenic, fluorine, and selenium from indoor burning of Chinese coal. *Rev Environ Contam Toxicol* 189:89–106.
- Stranges S, Marshall JR, Natarajan R, Donahue RP, Trevisan M, Combs GF, et al. 2007. Effects of long-term selenium supplementation on the incidence of type 2 diabetes: a randomized trial. *Ann Intern Med* 147(4):217–223.
- Stranges S, Navas-Acien A, Rayman MP, Guallar E. 2010. Selenium status and cardiometabolic health: state of the evidence. *Nutr Metab Cardiovasc Dis* 20(10):754–760.
- Sutter ME, Thomas JD, Brown J, Morgan B. 2008. Selenium toxicity: a case of selenosis caused by a nutritional supplement. *Ann Intern Med* 148(12):970–971.
- U.S. Environmental Protection Agency. 2002. Selenium and Compounds (CASRN 7782-49-2). Available: <http://www.epa.gov/iris/subst/0472.htm> [accessed 3 March 2011].
- Valera B, Dewailly É, Poirier P. 2009. Environmental mercury exposure and blood pressure among Nunavik Inuit adults. *Hypertension* 54(5):981–986.
- Vinceti M, Bonvicini F, Rothman KJ, Vescovi L, Wang F. 2010. The relation between amyotrophic lateral sclerosis and inorganic selenium in drinking water: a population-based case-control study. *Environmental Health* 9:77; doi:10.1186/1476-069X-9-77 [6 December 2010].
- Yang G, Wang S, Zhou R, Sun S. 1983. Endemic selenium intoxication of humans in China. *Am J Clin Nutr* 37(5):872–881.

ERRATUM

The February 2011 article “Avoiding Health Pitfalls of Home Energy-Efficiency Retrofits” [*Environ Health Perspect* 119:A76–A79 (2011)] incorrectly listed 1,2,3-trichloropropane (TCP) as a flame retardant used in spray polyurethane foams. The correct compound should have been tris(1-chloro-2-propyl) phosphate (TCPP).

EHP regrets the error.

for Educators



Locate lessons by topics

It's healthy, it's smart, and it's free.

www.ehponline.org/education